arXiv:1210.2812v2 [quant-ph] 29 Nov 2012

POLYNOMIAL CONSTRAINTS ON REPRESENTING ENTANGLED QUBITS AS MATRIX PRODUCT STATES

ANDREW CRITCH AND JASON MORTON

Abstract. We quantify the representational power of matrix product states (MPS) for entangled qubit systems by giving polynomial expressions in a pure quantum state's amplitudes which hold if and only if the state is a translation invariant matrix product state or a limit of such states. For systems with few qubits, we give these equations explicitly, considering both periodic and open boundary conditions. Using the classical theory of trace varieties and trace algebras, we explain the relationship between MPS and hidden Markov models and exploit this relationship to derive useful parameterizations of MPS. We make four conjectures on the identifiability of MPS parameters.

Matrix product states (MPS) provide a useful model of 1-D quantum spin systems which approximate the ground states of gapped local Hamiltonians [16]. Accordingly the problem of classifying phases of matter for such chains has been reduced to understanding equivalence classes (such as under LU operations) in the space of quantum states representable as matrix product states [17, 8, 5].

With periodic or open boundary conditions, we describe the closure of this space of states representable by translation invariant binary MPS as an algebraic variety. Our description is given as an ideal of polynomials in the state's amplitudes that vanish if and only if the state is a limit of MPS with N spins and $D = d = 2$ dimensional virtual and physical bonds. In small cases our description is complete. In Section 1, we exhibit a polynomial which vanishes on a pure state if and only if it is a limit of binary translation invariant, periodic boundary MPS with $N = 4$, and a set of 30 polynomials which vanish when $N = 5$. We also obtain many linear equations which are satisfied for N up to 12. In Section 2, Theorem 7 gives an analogous result for MPS with open boundary conditions and $N = 3$. Finally we examine cases where $N \gg 0$.

Matrix product states bear a close relationship to probabilistic graphical models known as a *hidden Markov models* (HMM)[4]. In Section 3, we make this relationship precise by modifying the parametrization of HMM to obtain MPS. We review the invariant theory of trace identities and trace varieties that has been used to study HMM[1], and how these results apply to varieties of MPS. In particular we obtain a nice parametrization for translation invariant binary MPS with periodic boundary conditions. Finally in Section 4 we suggest other such relationships between probabilistic graphical models and tensor network state models. Our results are complimentary to the connection between invariant theory and diagrammatic representations explored in [3] and the approaches to quantum state tomography for MPS developed in [2, 9].

1. Representability by translation invariant matrix product states

First consider a translation-invariant matrix product state with *periodic boundary* conditions. Suppose the inner (virtual) bond dimension is D, the outer (physical) bond dimension is d, and there are N spins. Fix $D \times D$ complex parameter matrices $A_0, \dots, A_{d-1}$, defining the same $D \times D \times d$ parameter tensor at each site. This defines the tensor network state, for $i_j \in \{0, \dots, d-1\}$,

$$\Psi = \sum_{i_1,\dots,i_N} \operatorname{tr}(A_{i_1} \cdots A_{i_N}) |i_1 i_2 \cdots i_N\rangle. \tag{1}$$

Question 1. *Fixing virtual and physical bond dimension, which states are matrix product states?*

Including states which are limits of MPS, a precise answer to this question could be given as a constructive description of the set of polynomials f in the coefficients of Ψ such that $f(\psi_{i_1,\dots i_N}) = 0$ if and only if ψ is a limit of MPS. This would describe the (closure of the) set of MPS as an algebraic variety. See [6] for background on varieties.

Such a description is possible because of the way MPS are defined. Each coefficient $\psi_{i_1,\dots,i_N}$ is a polynomial function of the parameters a_{rst} in the $D \times D \times d$ tensor A. Thus (1) defines a regular map $\Psi : \mathbb{C}^{D^2 d} \to \mathbb{C}^{d^N}$, whose image we denote by $\mathrm{PB}(D, d, N)$, the set of tensors representable by translation-invariant matrix product states with periodic boundary conditions. Its closure $\overline{\mathrm{PB}}(D, d, N)$ in either the Zariski or classical topology is an irreducible algebraic variety consisting of those tensors which can be approximated *arbitrarily well* by MPS. We can thus refine Question 1 as follows.

Question 2. *Fixing, D, d, and N, what polynomial relations must the coefficients of a matrix product state satisfy: what is the defining ideal of $\overline{\mathrm{PB}}(D, d, N)$?*



FIGURE 1. Translation-invariant MPS with periodic boundary.

We primarily examine the fully binary case $D = d = 2$. The invariance of trace under cyclic permutations of the matrices $A_{i_1}, \ldots, A_{i_N}$ means we can immediately restrict to the subspace spanned by *binary necklaces* (equivalence classes of binary strings under cyclic permutation). For $N = 3$ physical legs, this is the coordinate subspace $(\psi_{000} : \psi_{100} : \psi_{110} : \psi_{111})$ and all three-qubit states with cyclic symmetry are matrix product states. For $N = 4$ it is the six-dimensional coordinate subspace $(\psi_{0000} : \psi_{1000} : \psi_{1100} : \psi_{1010} : \psi_{1110} : \psi_{1111})$ and not all states are MPS (Theorem 3). In the $N = 5$ case the 8 equivalence classes of coefficients under cyclic permutation are ψ_{00000}, ψ_{10000}, ψ_{11000}, ψ_{11100}, ψ_{11110}, ψ_{11111}, ψ_{10100}, and ψ_{11010}. For $N = 6, \ldots, 15$ the



FIGURE 2. The eight binary necklaces for $N = 5$.

dimensions of this "necklace space" are 14, 20, 36, 60, 108, 188, 352, 632, 1182, and 2192. In general there are

$$n_d(N) = \frac{1}{N} \sum_{\ell | N} \varphi(\ell) d^{N/\ell}$$

d-ary necklaces of length N, where φ is Euler's totient function. Thus translation invariant MPS with periodic boundary of length N and physical bond dimension d live in a linear space isomorphic to $\mathbb{C}^{n_d(N)}$.

Naïvely we have 8 parameters in our 2×2×2 tensor A, but on each virtual bond we can apply a gauge transformation $P(\,\cdot\,)P^{-1}$ for $P \in SL_2$ without changing the state[13]. Since SL_2 is 3-dimensional, we expect $\overline{\mathrm{PB}}(2, 2, 3)$ to be 5-dimensional. Counting this way, our *expected dimension* of $\overline{\mathrm{PB}}(D, d, N)$ is $\min\{D^2(d-1) + 1, n_d(N)\}$. We expect $\overline{\mathrm{PB}}(D, d, N)$ to be a hypersurface when this equals $n_d(N)$, which happens first when $(D, d, N) = (2, 2, 4)$. In this case our expectation holds:

Theorem 3. *A four-qubit state Ψ is a limit of binary periodic translation invariant MPS with $N = 4$ if and only if the following irreducible* *polynomial vanishes:*

$$\begin{aligned}
&\psi_{1010}^2\psi_{1100}^4 - 2\psi_{1100}^6 - 8\psi_{1000}\psi_{1010}\psi_{1100}^3\psi_{1110} + 12\psi_{1000}\psi_{1100}^4\psi_{1110}\\
&- 4\psi_{1000}^2\psi_{1010}^2\psi_{1110}^2 + 2\psi_{0000}\psi_{1010}^3\psi_{1110}^2 + 16\psi_{1000}^2\psi_{1010}\psi_{1100}\psi_{1110}^2\\
&- 4\psi_{0000}\psi_{1010}^2\psi_{1100}\psi_{1110}^2 - 16\psi_{1000}^2\psi_{1100}^2\psi_{1110}^2 + 4\psi_{0000}\psi_{1010}\psi_{1100}^2\psi_{1110}^2\\
&- 4\psi_{0000}\psi_{1100}^3\psi_{1110}^2 - 4\psi_{0000}\psi_{1000}\psi_{1010}\psi_{1110}^3 + 8\psi_{0000}\psi_{1000}\psi_{1100}\psi_{1110}^3\\
&- \psi_{0000}^2\psi_{1110}^4 + 2\psi_{1000}^2\psi_{1010}^3\psi_{1111} - \psi_{0000}\psi_{1010}^4\psi_{1111} - 4\psi_{1000}^2\psi_{1010}^2\psi_{1100}\psi_{1111}\\
&+ 4\psi_{1000}^2\psi_{1010}\psi_{1100}^2\psi_{1111} + 2\psi_{0000}\psi_{1010}^2\psi_{1100}^2\psi_{1111} - 4\psi_{1000}^2\psi_{1100}^3\psi_{1111}\\
&+ \psi_{0000}\psi_{1100}^4\psi_{1111} - 4\psi_{1000}^3\psi_{1010}\psi_{1110}\psi_{1111} + 4\psi_{0000}\psi_{1000}\psi_{1010}^2\psi_{1110}\psi_{1111}\\
&+ 8\psi_{1000}^3\psi_{1100}\psi_{1110}\psi_{1111} - 8\psi_{0000}\psi_{1000}\psi_{1010}\psi_{1100}\psi_{1110}\psi_{1111}\\
&- 2\psi_{0000}\psi_{1000}^2\psi_{1110}^2\psi_{1111} + 2\psi_{0000}^2\psi_{1010}\psi_{1110}^2\psi_{1111} - \psi_{1000}^4\psi_{1111}^2\\
&+ 2\psi_{0000}\psi_{1000}^2\psi_{1010}\psi_{1111}^2 - \psi_{0000}^2\psi_{1010}^2\psi_{1111}^2.
\end{aligned}$$

Hence, up to closure, the the set $\mathrm{PB}(2,2,4)$ of tensors that can be represented in the form (1) where A_0 and A_1 are arbitrary 2×2 matrices, is a sextic hypersurface in the space of $2 \times 2 \times 2 \times 2$ tensors invariant under cyclic permutations of the indices. The 30-term hypersurface equation was found using a parametrization of the matrices that is similar to the birational parametrization of binary hidden Markov models given in [1].

An example of a pure state on four qubits on which the polynomial f of Theorem 3 is nonvanishing, and so cannot be arbitrarily well approximated by such a matrix product state, is given by letting $\psi_{1010} = \psi_{1110} = -1/4$ and $\psi_{0000} = \psi_{1000} = \psi_{1100} = \psi_{1111} = 1/4$. In this example, $f(\Psi) = 2^{-5}$, which is the maximal value of $f(\Psi)$ attained on corners of the 6-D hypercube.

The other cases with $N \leq 15$ when we expect $\overline{\mathrm{PB}}$ to be a hypersurface are when $(D, d, N) = (2, 4, 6)$, $(3, 3, 7)$, $(5, 15, 12)$, $(3, 71, 13)$, and $(2, 296, 14)$. In general, we will need many more polynomials to define the space of matrix product states as their zero locus. As an example, consider $\overline{\mathrm{PB}}(2, 2, 5)$, which we expect to be a five-dimensional variety in the necklace space $\mathbb{C}^8 = \mathbb{C}^{N_2(5)}$.

Theorem 4. *Any homogeneous minimal generating set for the ideal of* $\overline{\mathrm{PB}}(2, 2, 5)$ *must contain exactly 3 quartic and 27 sextic polynomials, possibly some higher degree polynomials, but none of degree* 1, 2, 3, *or* 5.

Proof. Using the bi-grading of Theorem 5, we decompose the ideal I into vector spaces $I_{r,s}$. For each (r, s) with $\frac{1}{5}(r+s) \leq 6$, we select a large number of parameter values $\widehat{A}$ at random, and use Gaussian elimination to compute a basis for the vector space $\widehat{I_{r,s}}$ of polynomials vanishing at their images $\Psi(\widehat{A})$, which is certain to contain $I_{r,s}$. We then substitute indeterminate entries for A symbolically into the polynomials to ensure that they lie in $I_{r,s}$, yielding a bihomogeneous basis for I in total degree ≤ 6. □

This is interesting, because the variety only has codimension 3, but requires *at least* 30 equations to cut it out ideal-theoretically. Such a collection of 3 quartics and 27 quadrics was found and verified symbolically. Exact numerical tests (intersection with random hyperplanes) indicate that the top dimensional component of the ideal they generate is reduced and irreducible of dimension 5, and is therefore equal to $\overline{\mathrm{PB}}(2,2,5)$.

1.1. **Homogeneity and GL_d-invariance.**

Note that the equation of Theorem 3 is homogeneous of degree 6, and every monomial has the same total number of 1s appearing in its subscripts. Every MPS variety will be homogeneous in such a grading:

Proposition 5. *For any D, d, N, the space of translation-invariant MPS limits with periodic boundary conditions is cut out by polynomials in which each monomial has the same total number of* 0*s,* 1*s, …* $(d-1)$*s appearing in its subscripts.*

Proof. In fact we claim that the ideal of $\overline{\mathrm{PB}}(D,d,N)$ is $\mathbb{Z}^d$-homogeneous with respect to d different $\mathbb{Z}$-gradings $\deg_i$ for $0 \leq i \leq d-1$, where $\deg_i(\Psi_J)$ is the number of occurrences of i in J. Since $\deg(\psi_J) := \frac{1}{n}\sum_{i=0}^{N-1} \deg_i(\psi_J) = 1$, $\overline{\mathrm{PB}}(D,d,N)$ is also homogeneous in the standard grading.

The usual parametrization Ψ, where $A_0, \ldots, A_{d-1}$ have generic entries, is $\mathbb{Z}^d$-homogeneous with respect to the grading above along with letting $\deg_i(A_j) = 1$ when $i = j$ and 0 when $i \neq j$. Being a homogeneous map, its kernel, the defining ideal of $\overline{\mathrm{PB}}(D,d,N)$, is homogeneous in each of these gradings as well. □

In fact, the variety is homogeneous in a stronger sense because of an action of GL_d on the parameter space of Ψ. In the example above, the action is given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}(A_0, A_1) = (aA_0 + bA_1, cA_1 + dA_1)$$

which descends to an action on Ψ by

$$\begin{pmatrix} g_{00} & g_{01} \\ g_{10} & g_{11} \end{pmatrix} \cdot \psi_{ijkl} = \sum_{pqrs} g_{ip} g_{jq} g_k r g_{ls} \psi_{pqrs}.$$

The embedding $(\mathbb{C}^*)^d \subset GL_d$ as diagonal matrices gives rise to the $\mathbb{Z}^d$ homogeneity of the proposition above.

1.2. **Linear invariants and reflection symmetry.** There are additional symmetries peculiar to the case $D = d = 2$. For a generic pair of 2×2 matrices A_0, A_1, there is a one-dimensional family of matrices $P \in SL_2$ such that $P^{-1} A_i P$ are symmetric. Thus, a generic point $\Psi \in \mathrm{PB}(2, 2, N)$ can be written as $\Psi(A_0, A_1)$ with $A_i^T = A_i$, and then $\Psi_J = \mathrm{tr}(\prod_{j \in J} A_j) = \mathrm{tr}((\prod_{j \in J} A_j)^T) = \mathrm{tr}(\prod_{j \in \mathrm{reverse}(J)} A_j) = \Psi_{\mathrm{reverse}(J)}$. This implies

Proposition 6. *If an N-qubit state Ψ is a limit of binary periodic translation invariant matrix product states, then it has reflection symmetry: $\psi_J = \psi_{\mathrm{reverse}(J)}$ for all J.*

For $N \geq 6$, N-bit strings can be equivalent under reflection but not cyclic permutation, so then $\mathrm{PB}(2, 2, N)$ admits additional linear invariants, i.e. linear polynomials vanishing on the model. For $N = 6, 7$ these are:

$$\begin{aligned} \mathrm{PB}(2,2,6): &\quad \psi_{110100} - \psi_{110010} \\ \mathrm{PB}(2,2,7): &\quad \psi_{1110100} - \psi_{1110010} \text{ and } \psi_{1101000} - \psi_{1100010} \end{aligned}$$

For small N we can find *all* the linear invariants of $\mathrm{PB}(2, 2, N)$ using the bigrading of Theorem 5 as in the proof of Theorem 4. Modulo the cyclic and reflection invariants, there are no further linear invariants for $N \leq 7$, but $\mathrm{PB}(2, 2, 8)$ has a single "non-trivial" linear invariant,

$$\psi_{11010010} + \psi_{11001100} - \psi_{11001010} + \psi_{11101000} - \psi_{11011000} - \psi_{11100100}.$$

For $N = 9$, 10, 11, and 12, $\mathrm{PB}(2, 2, N)$ admits 6, 17, 44, and 106 such non-trivial invariants, in each case unique up to change of basis on the vector space they generate.

2. MPS WITH OPEN BOUNDARY CONDITIONS

We now consider matrix product states with open boundary conditions, which are even more similar to hidden Markov models than the periodic version. Here the state is determined by two boundary state vectors $b_0, b_1 \in \mathbb{C}^D$, along with the $D \times D$ parameter matrices $A_0, \ldots, A_{d-1}$ of the MPS, by

$$\Psi = \sum_{i_1,\ldots,i_N} b_0^T A_{i_1} \cdots A_{i_N} b_1 |i_1 i_2 \cdots i_N\rangle \tag{2}$$

$$= \sum_{i_1,\ldots,i_N} \operatorname{tr}(B A_{i_1} \cdots A_{i_N}) |i_1 i_2 \cdots i_N\rangle \tag{3}$$

where $B = b_1 b_0^T$ is a rank 1 matrix. We denote the set of states obtainable in this way by $\mathrm{OB}(D, d, N)$, and its closure (Zariski or classical) by $\overline{\mathrm{OB}}(D, d, N)$. We do not have the cyclic symmetries of the PB model here, so we consider $\overline{\mathrm{OB}}(D, d, N)$ as a subvariety of $\mathbb{C}^{d^N}$. If the A_i and b_0^T have non-negative entries with row sums equal to 1, and b_1 is a vector of 1's, then (2) is exactly the Baum formula for HMM, so in fact the model $\mathrm{HMM}(D, d, N)$ studied in [1] is contained in $\mathrm{PB}(D, d, N)$.

The expression (3) for Ψ is invariant under the action of SL_D on the A_i and B by simultaneous conjugation. Thus, we may assume B is in Jordan normal form, i.e. a matrix of all zeroes except possibly in the top left corner. As well, the map $(B, A_1, \ldots, A_d) \mapsto (t^{-N}B, tA_1, \ldots, tA_d)$ preserves Ψ, so discarding the case $B = 0$ (which will not change $\overline{\mathrm{OB}}$) we can assume that the top left entry of B is 1. Thus Ψ is determined by dD^2 parameters, the entries of the A_i. In particular, $\overline{\mathrm{OB}}(2, 2, 3)$ is parametrized by (a dominant map from) 8 parameters, and lives in an 8-dimensional space. This parametrization still turns out still to be degenerate:

Theorem 7. *A three-qubit state Ψ is a limit of $N = 3$ binary translation invariant MPS with open boundary conditions if and only if the following 22-term quartic polynomial vanishes:*

$$\begin{aligned}
&\psi_{011}^2\psi_{100}^2 - \psi_{001}\psi_{011}\psi_{100}\psi_{101} - \psi_{010}\psi_{011}\psi_{100}\psi_{101} + \psi_{000}\psi_{011}\psi_{101}^2 \\
&+ \psi_{001}\psi_{010}\psi_{011}\psi_{110} - \psi_{000}\psi_{011}^2\psi_{110} - \psi_{010}\psi_{011}\psi_{100}\psi_{110} \\
&+ \psi_{001}\psi_{010}\psi_{101}\psi_{110} + \psi_{001}\psi_{100}\psi_{101}\psi_{110} - \psi_{000}\psi_{101}^2\psi_{110} - \psi_{001}^2\psi_{110}^2 \\
&+ \psi_{000}\psi_{011}\psi_{110}^2 - \psi_{001}\psi_{010}^2\psi_{111} + \psi_{000}\psi_{010}\psi_{011}\psi_{111} + \psi_{001}^2\psi_{100}\psi_{111} \\
&+ \psi_{010}^2\psi_{100}\psi_{111} - \psi_{000}\psi_{011}\psi_{100}\psi_{111} - \psi_{001}\psi_{100}^2\psi_{111} - \psi_{000}\psi_{001}\psi_{101}\psi_{111} \\
&+ \psi_{000}\psi_{100}\psi_{101}\psi_{111} + \psi_{000}\psi_{001}\psi_{110}\psi_{111} - \psi_{000}\psi_{010}\psi_{110}\psi_{111}
\end{aligned}$$

That is, the variety $\overline{\mathrm{OB}}(2, 2, 3)$ is a quartic hypersurface in $\mathbb{C}^8$ cut out by the polynomial above. This polynomial previously appeared in the context of the HMM[12]*.*

Proof. The map Ψ and its image are homogeneous in the same grading as described in Theorem 5, which we can use as in the proof of

Theorem 4 to search for low degree polynomials vanishing on the variety. When $(D, d, N) = (2, 2, 3)$ the quartic from the theorem appears in this search. The quartic is prime, and therefore defines a 7-dimensional irreducible hypersurface in $\mathbb{C}^8$. On the other hand, the Jacobian of the map Ψ at a random point, e.g. the point where A_0, A_1 have entries $1, 2, 3, 4, 5, 6, 7, 8$ in that order, has rank 7. Therefore $\overline{\mathrm{OB}}(2, 2, 3)$ is of dimension at least 7, and contained in the quartic hypersurface above, so they must be equal. □

From Theorem 7, we can derive conditions on $\mathrm{OB}(2, 2, N)$ for $N \geq 4$ as well. There is a *marginalization map* from $\mathrm{OB}(2, 2, N)$ to $\mathrm{OB}(2, 2, 3)$ given by $\Psi_I \mapsto \sum_{|J|=N-3} \Psi_{IJ}$ for each I of length 3, which commutes with the assignment $b_1 \mapsto \sum_{j_3 \ldots j_N} A_{j_3} \cdots A_{j_N} b_1$. In fact there are $N - 2$ such marginalization maps, each given by choosing 3 consecutive indices I to marginalize to (summing over the remaining indices J). By composing these maps with the quartic polynomial above, we can obtain $N - 2$ quartic polynomials vanishing on $\mathrm{OB}(2, 2, N)$.

By analogy to the case of hidden Markov models discussed in the next section, we make the following

Conjecture 8. *For $N \geq 4$, a generic N-qubit state can be recovered from its marginalization to any three consecutive states. That is, each marginalization map $\overline{\mathrm{OB}}(2, 2, N) \to \overline{\mathrm{OB}}(2, 2, 3)$ is a birational equivalence of varieties.*

The analogous statement with $\overline{\mathrm{HMM}}$ in place of $\overline{\mathrm{OB}}$ is shown to be true in [1].

Conjecture 9. *A generic N-qubit (D=d=2) translation invariant matrix product state Ψ with open boundary conditions is determined up to phase by a reduced density operator which traces out all but a chain of three adjacent states, but no fewer.*

When the three adjacent states are qubits 1, 2, and 3 (the first three legs of the diagram), this amounts to saying that the group S^1 of unit-modulus complex numbers acts transitively on generic fibres of the real-algebraic map

$$\Psi \mapsto \left(\sum_{i_4, \ldots, i_N} \Psi_{j_1 j_2 j_3 i_4 \ldots i_N} \Psi^{\dagger}_{k_1 k_2 k_3 i_4 \ldots i_N} \right)_{j_1 j_2 j_3 k_1 k_2 k_3}$$

when restricted to $\mathrm{OB}(2, 2, N)$. Here the right hand side denotes an order 6 tensor with indices j_1, j_2, j_3, k_1, k_2, k_3, and $\Psi^{\dagger}$ denotes complex conjugation.



FIGURE 3. Parameterization of an MPS model as a complex HMM using complex E and T matrices with all row sums equal to $z \in \mathbb{C}$ and copy dot (comultiplication) tensor (circle). Contraction of a region of the tensor network enclosed by a dashed line yields an A tensor.

Conjecture 10. *A generic N-qubit (D=d=2) periodic translation invariant matrix product state Ψ is determined up to phase by a reduced density operator which traces out all but a chain of four adjacent states, but no fewer.*

Similarly, this amounts to saying that S^1 acts transitively on generic fibres of the map

$$\Psi \mapsto \left(\sum_{i_5,\ldots,i_N} \Psi_{j_1 j_2 j_3 j_4 i_5 \ldots i_N} \Psi^{\dagger}_{k_1 k_2 k_3 k_4 i_5 \ldots i_N} \right)_{j_1 j_2 j_3 j_4 k_1 k_2 k_3 k_4}$$

when restricted to $\mathrm{PB}(2, 2, N)$.

3. MATRIX PRODUCT STATES AS COMPLEX VALUED HIDDEN MARKOV MODELS

We now explain how the polynomial in Theorem 3 was obtained, and connect the classical *hidden Markov model* and matrix product states through a reparametrizing rational map. The parametrization of the state Ψ is analogous to that of the moment tensor of a binary hidden Markov model used in [1] for symbolic computations.

Let T be a 2×2 *transition* matrix and E a 2×2 *emission* matrix. For a (classical) hidden Markov model, T and E are nonnegative stochastic matrices (their rows sum to one), representing a four-dimensional parameter space. For PB, T and E will be complex with row sums all equal to some constant $z \in \mathbb{C}$, so they form parameter space isomorphic to $\mathbb{C}^5$. We parametrize the A_i in terms of (T, E) by

$$A_0 = T, \qquad A_1 = \begin{pmatrix} e_{01} & 0 \\ 0 & e_{11} \end{pmatrix} \begin{pmatrix} t_{00} & t_{01} \\ t_{10} & t_{11} \end{pmatrix}.$$

This is shown in Figure 3; grouping and contracting the E, T, and copy dot tensors into an A tensor yields a dense parameterization of an MPS as depicted in Figure 1. We then parameterize E and T with the five parameters u, v_0, b, c_0, z by setting

$$\begin{aligned} E &= \begin{pmatrix} z-u+v_0 & u-v_0 \\ z-u-v_0 & u+v_0 \end{pmatrix} \text{ and} \\ T &= \begin{pmatrix} z+u-c_0 & z-b+c_0 \\ z-b-c_0 & z+b+c_0 \end{pmatrix}. \end{aligned}$$

Composing these formulae with the map $(A_0, A_1) \mapsto \Psi$ yields a restricted parametrization $\rho_N : \mathbb{C}^5 \to \mathbb{C}^{2^N}$, whose image lies inside $\mathrm{PB}(2,2,N)$.

Proposition 11. *The variety* $\overline{\mathrm{PB}}(2,2,N)$ *is at most* 5*-dimensional, and the image of our restricted parametrization* ρ_N *is dense in it.*

Proof. Suppose $\Psi = \Psi(A_0, A_1)$ for A_0, A_1 generic. First, we will transform the A_i by simultaneous conjugation with an element P of SL_2 to a new pair of matrices A_0', A_1' such that A_0' has equal row sums and $A_1' = DA_0'$ for a diagonal matrix D. Generically, A_0 is invertible, and we can diagonalize the matrix $A_1A_0^{-1}$, so we write $U^{-1}A_1A_0^{-1}U = D_0$, and then $U^{-1}A_1U = DU^{-1}A_0U$. Next we find another diagonal matrix $D_1 \in SL_2$ such that $D_1^{-1}U^{-1}A_0UD_1$ has equal row sums. Then let $P = UD_1$ and $A_i' = D_1^{-1}U^{-1}A_iUD_1$, and we are done with our transformation. Now $\Psi = \Psi(A_0', A_1')$ since simultaneous conjugation does not change trace products. But now letting z be the common row sum of A_0', we can solve linearly for u, v_0, b, and c_0 to obtain $\Psi = \rho(u, v_0, b, c_0, z)$. □

In fact we know from exact computations in Macaulay2[7] that the dimension $\dim \mathrm{PB}(2,2,N) = 5$ for $4 \leq N \leq 100$. This is proven by checking that the Jacobian of ρ attains rank 5 at some point with randomly chosen integer coordinates, giving a lower bound of 5 on the dimension of its image.

When parametrized using ρ, there are sufficiently few parameters and the entries of Ψ are sufficiently short expressions that Macaulay2 is also able to compute the exact kernel of the parametrization, i.e. defining equations for the model. It is by this method that we obtain the hypersurface equation of Theorem 3 as the only ideal generator for $\mathrm{PB}(2,2,4)$.

3.1. **Identifying parameters of MPS.** Determining the parameters of an MPS is related to *quantum state tomography*, and represents a quantum analog to the identifiability problem in statistics. The extent to which the parameters can be identified can be addressed algebraically.

Given $D \times D$ matrices $A_1 \dots A_d$ with indeterminate entries, we write $\mathcal{C}_{D,d}$ for the algebra of polynomial expressions in their entries that are invariant under simultaneous conjugation of the matrices by GL_2.

Sibirskii[15], Leron[10], and Procesi[14] showed that the algebra $\mathcal{C}_{D,d}$ is generated by the traces of products $tr(A_{i_0} \cdots A_{i_n})$ as $n \geq 0$ varies. For this reason, $\mathcal{C}_{D,d}$ is called a *trace algebra.* Its spectrum, $\operatorname{Spec} \mathcal{C}_{D,d}$, is a *trace variety.* Since the coordinate ring of $\overline{\mathrm{PB}}(D,d,N)$ is a subring of $\mathcal{C}_{D,d}$, we have a map $\operatorname{Spec} \mathcal{C}_{D,d} \to \mathbb{C}^{d^N}$ parametrizing a dense open subset of $\overline{\mathrm{PB}}(D,d,N)$.

In the case $D = 2$, Sibirskii showed further that the trace algebra $\mathcal{C}_{2,d}$ is minimally generated by the elements $tr(A_i)$ and $tr(A_i^2)$ for $1 \leq i \leq d$, $tr(A_iA_j)$ for $1 \leq i < j \leq d$, and $tr(A_iA_jA_k)$ for $1 \leq i < j < k \leq d$.

For $d = 1 \dots 6$, the number of such generators is $2, 5, 10, 18, 30, 47$. In particular, when $d = 2$, the number of generators equals the transcendence degree of the ring, $5 = 8 - 3$. This means $\operatorname{Spec} C_{2,2}$ is isomorphic to $\mathbb{C}^5$, yielding for each N a dominant parametrization $\phi_N : \mathbb{C}^5 \to \overline{\mathrm{PB}}(2,2,N)$. Gröbner bases for randomly chosen fibers indicate that for $N = 4 \dots 10$, the map ϕ_N is generically k-to-one, where $k = 8, 5, 6, 7, 8, 9, 10$, respectively. Continuing this sequence suggests the following.

Conjecture 12. *Using the trace parameterization ϕ_N, for $N \geq 5$, almost every periodic boundary MPS has exactly N choices of parameters that yield it.*

In other words, for $N \geq 5$, the parametrization $\phi_N : \mathbb{C}^5 \simeq \operatorname{Spec} C_{2,2} \to \overline{\mathrm{PB}}(2,2,N)$ is generically N-to-1. Generically, the points of $\operatorname{Spec} C_{2,2}$ are in bijection with the SL_2-orbits of the tensors A. The conjecture implies that, up to the action of SL_2, the parameters of a binary, $D = d = 2$ translation invariant matrix product state with periodic boundary are algebraically identifiable from its entries.

4. CONCLUSION

A conjectured dictionary between tensor network state models and classical probabilistic graphical models was presented in [11]. In this dictionary, matrix product states correspond to hidden Markov models, the density matrix renormalization group (DMRG) algorithm to the forward-backward algorithm, tree tensor networks to general Markov models, projected entangled pair states (PEPS) to Markov or conditional random fields, and the multi-scale entanglement renormalization ansatz (MERA) loosely to deep belief networks. In this work we formalize the first of these correspondences and use it to algebraically characterize quantum states representable by MPS and study their identifiability.

Acknowledgments. AC and JM were supported in part by DARPA under awards FA8650-10-C-7020 and N66001-10-1-4040 respectively. We would like to thank J. Biamonte, J. Eisert, B. Sturmfels, F. Vaccarino, F. Verstraete, and G. Vidal for helpful discussions.

References

[1] Andrew J. Critch. Binary hidden Markov models and varieties. arXiv:1206.0500, 2012.

[2] T. Baumgratz, D. Gross, M. Cramer, and M.B. Plenio. Scalable reconstruction of density matrices. *arXiv preprint arXiv:1207.0358*, 2012.

[3] J. Biamonte, V. Bergholm, and M. Lanzagorta. Invariant theory for matrix product states. arXiv:1209.0631, 2012.

[4] N. Bray and J. Morton. Equations defining hidden Markov models. In L. Pachter and B. Sturmfels, editors, *Algebraic Statistics for Computational Bio.*, chapter 11, pages 237–249. Camb. Univ. Press, 2005.

[5] X. Chen, Z.C. Gu, and X.G. Wen. Classification of gapped symmetric phases in one-dimensional spin systems. *Physical Review B*, 83(3):035107, 2011.

[6] D Cox, J Little, and D O'Shea. *Ideals, varieties, and algorithms.* Springer-Verlag, New York, second edition, 1997.

[7] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/.

[8] M. B. Hastings. Entropy and entanglement in quantum ground states. *Phys. Rev. B*, 76:035114, 2007.

[9] R. Hübener, A. Mari, and J. Eisert. Wick's theorem for matrix product states. *arXiv preprint arXiv:1207.6537*, 2012.

[10] U. Leron. Trace identities and polynomial identities of $n \times n$ matrices. *J. Algebra*, 42:369–377, 1976.

[11] J. Morton. Tensor networks in algebraic geometry and statistics. Lecture at Networking Tensor Networks, Centro de Ciencias de Benasque Pedro Pascual, Benasque, Spain, May 2012.

[12] L. Pachter and B. Sturmfels. Tropical geometry of statistical models. *PNAS*, 101(46):16132–16137, 2004.

[13] D. Perez-Garcia, F. Verstraete, MM Wolf, and JI Cirac. Matrix product state representations. *Quantum Information & Computation*, 7(5):401–430, 2007.

[14] C. Procesi. The invariant theory of $n \times n$ matrices. *Adv. Math.*, 19:306–381, 1976.

[15] K.S. Sibirskii. Algebraic invariants for a set of matrices. *Siberian Mathematical Journal*, 9:115–124, 1968.

[16] F. Verstraete and J.I. Cirac. Matrix product states represent ground states faithfully. *Physical Review B*, 73(9):094423, 2006.

[17] F. Verstraete, J.I. Cirac, JI Latorre, E. Rico, and M.M. Wolf. Renormalization-group transformations on quantum states. *Physical review letters*, 94(14):140601, 2005.